

Existence of Positive Radial Solutions for Elliptic Systems

Ruyun Ma

Department of Mathematics, Northwest Normal University, Lanzhou 730070, Gansu, People's Republic of China; and Department of Mathematics, Lanzhou University, Lanzhou 730000, Gansu, People's Republic of China

etadata, citation and similar papers at core.ac.uk

1. INTRODUCTION

In this paper, we consider the existence of positive radial solutions of the elliptic system

$$\begin{aligned}\Delta u + a(|x|)f(u, v) &= 0, \\ \Delta v + b(|x|)g(u, v) &= 0, \quad R_1 < |x| < R_2, \quad x \in \mathbb{R}^N, \quad N \geq 2\end{aligned}\quad (1.1)$$

with one of the following sets of boundary conditions

$$u = v = 0 \quad \text{on } |x| = R_1, |x| = R_2, \quad (1.2)_a$$

$$u = v = 0 \quad \text{on } |x| = R_1, \quad \frac{\partial u}{\partial r} = \frac{\partial v}{\partial r} = 0 \quad \text{on } |x| = R_2, \quad (1.2)_b$$

$$\frac{\partial u}{\partial r} = \frac{\partial v}{\partial r} = 0 \quad \text{on } |x| = R_1, \quad u = v = 0 \quad \text{on } |x| = R_2, \quad (1.2)_c$$

where $f: \mathbb{R}^+ \times \mathbb{R}^+ \rightarrow \mathbb{R}^+$, $g: \mathbb{R}^+ \times \mathbb{R}^+ \rightarrow \mathbb{R}^+$ are continuous, $a: \mathbb{R}^+ \rightarrow \mathbb{R}^+$, $b: \mathbb{R}^+ \rightarrow \mathbb{R}^+$ are continuous functions, and $\mathbb{R}^+ = [0, \infty)$.

The existence and uniqueness of positive radial solutions for scalar equation

$$\Delta u + h(|x|)\tilde{f}(u) = 0, \quad R_1 \leq |x| \leq R_2, \quad x \in \mathbb{R}^N, \quad N \geq 2, \quad (1.3)$$

have been widely studied (see [1, 5–7]). In [7], Wang studied (1.3) with one of the following sets of boundary conditions

$$u = 0 \quad \text{on } |x| = R_1, |x| = R_2, \quad (1.4)_a$$

$$u = 0 \quad \text{on } |x| = R_1, \quad \frac{\partial u}{\partial r} = 0 \quad \text{on } |x| = R_2, \quad (1.4)_b$$

$$\frac{\partial u}{\partial r} = 0 \quad \text{on } |x| = R_1, \quad u = 0 \quad \text{on } |x| = R_2. \quad (1.4)_c$$

He proved the following results.

Assume $\tilde{f}: \mathbb{R}^+ \rightarrow \mathbb{R}^+$, $h: \mathbb{R}^+ \rightarrow \mathbb{R}^+$ are continuous, and h is not identically zero in any finite subinterval of $(0, \infty)$. If \tilde{f} satisfies

$$\lim_{u \rightarrow 0} \frac{f(u)}{u} = \infty, \quad \lim_{u \rightarrow \infty} \frac{f(u)}{u} = 0, \quad (1.5)$$

then (1.3)–(1.4) has a positive radial solution for any annulus $R_1 < |x| < R_2$.

The purpose of this paper is to give an extension of above-mentioned result of system (1.1)–(1.2). The proof of our main result (see Section 2) makes use of the following fixed point theorem.

FIXED POINT THEOREM [4]. *Let X be a Banach space, let K be a cone in X , let $0 < r < R$ be real numbers, let $D = \{x \in K : r \leq \|x\| \leq R\}$, and let $T: D \rightarrow K$ be a compact continuous operator such that*

- (i) $x \in D, \|x\| = R, Tx = \lambda x \Rightarrow \lambda \leq 1$;
- (ii) $x \in D, \|x\| = r, Tx = \lambda x \Rightarrow \lambda \geq 1$;
- (iii) $\inf_{\|x\|=r} \|Tx\| > 0$.

Then T has a fixed point in D .

2. MAIN RESULT

For $(x, y) \in \mathbb{R}^2$, denote $|(x, y)| = \max\{|x|, |y|\}$. The following conditions will be assumed in this section:

(H₁) $f: \mathbb{R}^+ \times \mathbb{R}^+ \rightarrow \mathbb{R}^+$, $g: \mathbb{R}^+ \times \mathbb{R}^+ \rightarrow \mathbb{R}^+$, $a: \mathbb{R}^+ \rightarrow \mathbb{R}^+$, $b: \mathbb{R}^+ \rightarrow \mathbb{R}^+$ are continuous.

(H₂) a and b are not identically zero in any finite subinterval of $(0, \infty)$.

(H₃)

$$\lim_{|(x, y)| \rightarrow 0} \frac{f(x, y)}{|(x, y)|} = \infty \quad (2.1)$$

and

$$\lim_{|(x, y)| \rightarrow 0} \frac{g(x, y)}{|(x, y)|} = \infty. \quad (2.2)$$

(H₄) Either

- (i) f and g are bounded functions, or
- (ii) g is bounded and f is unbounded and

$$\lim_{|(x, y)| \rightarrow \infty} \frac{f(x, y)}{|(x, y)|} = 0, \quad (2.3)$$

or

(iii) g is increasing in each variable for any value of the other variable, f is unbounded, and

$$\lim_{|(x, y)| \rightarrow \infty} \frac{f(x, y)}{|(x, y)|} = 0 = \lim_{|(x, y)| \rightarrow \infty} \frac{g(x, y)}{|(x, y)|}, \quad (2.4)$$

or

- (iv) f is bounded, g is unbounded, and

$$\lim_{|(x, y)| \rightarrow \infty} \frac{g(x, y)}{|(x, y)|} = 0,$$

or

(v) f is increasing in each variable for any value of the other variable, g is unbounded, and (2.4) holds.

THEOREM 1. Assume (H₁)–(H₄) hold. Then (1.1)–(1.2) has a positive radial solution for any annulus $R_1 < |x| < R_2$.

Remark 2. Theorem 1 is an extension of Theorem 1 of [7], which deals with the scalar equation (1.3) with the boundary conditions (1.4).

3. PRELIMINARIES

In view of the spherical symmetry of $g(|x|)$, we seek a positive radial solution $u = u(r), v = v(r)$ to (1.1). Therefore we write (1.1)–(1.2) in the

form

$$\begin{aligned}
 u''(r) + \frac{N-1}{r}u'(r) + a(r)f(u(r), v(r)) &= 0, \\
 v''(r) + \frac{N-1}{r}v'(r) + b(r)g(u(r), v(r)) &= 0, \quad R_1 < x < R_2, \quad (3.1) \\
 u(R_1) = v(R_1) = u(R_2) = v(R_2) &= 0, \quad (3.2)_a \\
 u(R_1) = v(R_1) = u'(R_2) = v'(R_2) &= 0, \quad (3.2)_b \\
 u'(R_1) = v'(R_1) = u(R_2) = v(R_2) &= 0, \quad (3.2)_c
 \end{aligned}$$

Let $s = -\int_r^{R_2}(1/t^{N-1})dt$ and $w(s) = u(r(s))$ and $z(s) = v(r(s))$, then (3.1)–(3.2) can be rewritten as

$$\begin{aligned}
 w''(s) + r^{2(N-1)}(s)a(r(s))f(w(s), z(s)) &= 0, \\
 z''(s) + r^{2(N-1)}(s)b(r(s))g(w(s), z(s)) &= 0, \quad m < s < 0, \\
 w(m) = z(m) = w(0) = z(0) &= 0, \\
 w(m) = z(m) = w'(0) = z'(0) &= 0, \\
 w'(m) = z'(m) = w(0) = z(0) &= 0,
 \end{aligned}$$

where $m = -\int_{R_1}^{R_2}(1/t^{N-1})dt$.

To be obvious, again let $t = (m - s)/m$ and $\varphi(t) = w(s)$ and $\psi(t) = z(s)$. Then (3.1)–(3.2) can also be written as

$$\begin{aligned}
 \varphi''(t) + A(t)f(\varphi(t), \psi(t)) &= 0, \\
 \psi''(t) + B(t)g(\varphi(t), \psi(t)) &= 0, \quad (3.3) \\
 \varphi(0) = \psi(0) = \varphi(1) = \psi(1) &= 0, \quad (3.4)_a \\
 \varphi(0) = \psi(0) = \varphi'(1) = \psi'(1) &= 0, \quad (3.4)_b \\
 \varphi'(0) = \psi'(0) = \varphi(1) = \psi(1) &= 0, \quad (3.4)_c
 \end{aligned}$$

where $A(t) = m^2 r^{2(N-1)}(m(1-t))a(r(m(1-t)))$, $B(t) = m^2 r^{2(N-1)}(m(1-t))b(r(m(1-t)))$. It is easy to check that $A(t)$ and $B(t)$ also satisfy (H_2) .

From now on, we concentrate on (3.3)–(3.4). Indeed, if we prove that there exists a positive solution to (3.3)–(3.4) for any $m \neq 0$, then (1.1)–(1.2) has a positive radial solution for any annulus.

In the proof of Theorem 1, we will use the following two lemmas.

LEMMA 3. If $f: \mathbb{R}^+ \rightarrow \mathbb{R}^+$ is a unbounded continuous function, then the set

$$\Sigma_f = \left\{ l \in \mathbb{R}^+ \left| \begin{array}{ll} \max_{[0, l; 0, l]} f(x, y) = f(l, \bar{y}) & \text{for some } \bar{y} \in [0, l], \text{ or} \\ \max_{[0, l; 0, l]} f(x, y) = f(\bar{x}, l) & \text{for some } \bar{x} \in [0, l] \end{array} \right. \right\} \quad (3.5)$$

is unbounded.

LEMMA 4. Let $h \in L^1(0, 1)$ and u be a solution of the equation $-u''(x) = h(x)$ with either $u(0) = u(1) = 0$ or $u(0) = u'(1) = 0$ or $u'(0) = u(1) = 0$. If $u \neq 0$, then $u(x) > 0$ for all $x \in (0, 1)$.

The proofs of Lemmas 3 and 4 follow from elementary arguments and are omitted.

4. PROOF OF THEOREM 1

In what follows X will denote the space of continuous functions $u: [0, 1] \rightarrow \mathbb{R}$ with the norm $\|u\| = \max_{0 \leq x \leq 1} |u(x)|$.

First consider (3.3)–(3.4)_a. Let

$$K_1 = \{ \varphi \in X : \varphi \text{ has graph concave down, } \varphi \geq 0, \varphi(0) = \varphi(1) = 0 \};$$

then for $\varphi \in K_1$, we have that $\varphi(t) \geq \frac{1}{4}\|\varphi\|$ for $t \in [\frac{1}{4}, \frac{3}{4}]$.

In $X \times X$ consider the maximum norm and define

$$T: K_1 \times K_1 \rightarrow K_1 \times K_1$$

given by

$$T(\varphi, \psi)(t) = \left(\int_0^1 G(t, s) A(s) f(\varphi(s), \psi(s)) ds, \right. \\ \left. \int_0^1 G(t, s) B(s) g(\varphi(s), \psi(s)) ds \right),$$

where

$$G(t, s) = \begin{cases} t(1-s), & t \leq s, \\ s(1-t), & t > s. \end{cases}$$

It is easy to see $G(t, s) \leq G(s, s)$.

Observe that, if we write

$$T(\varphi, \psi) = (\varphi_1, \psi_1),$$

then φ_1, ψ_1 are continuously differentiable on $[0, 1]$, twice differentiable on $(0, 1)$, and

$$\varphi_1'(t) = -\int_0^t sA(s)f(\varphi(s), \psi(s))ds + \int_t^1 (1-s)A(s)f(\varphi(s), \psi(s))ds,$$

$$\varphi_1''(t) = -A(s)f(\varphi(t), \psi(t)) \leq 0;$$

likewise for ψ_1 . This implies that T is well defined and the Arzela–Ascoli theorem implies that T is a compact mapping. It is easy to check that (3.3)–(3.4)_a is equivalent to the fixed point problem in $K_1 \times K_1$,

$$(\varphi, \psi) = T(\varphi, \psi).$$

From (H_3) , there is a $r > 0$ such that

$$\begin{aligned} f(x, y) &\geq M|(x, y)|, \\ g(x, y) &\geq M|(x, y)|, \end{aligned} \quad (4.1)$$

for any $(x, y) \in \mathbb{R}^+ \times \mathbb{R}^+ : 0 < |(x, y)| \leq r$, where the constant M satisfies

$$\frac{M}{4} \inf_{1/4 \leq t \leq 3/4} \min \left\{ \int_{1/4}^{3/4} G(t, s)A(s)ds, \int_{1/4}^{3/4} G(t, s)B(s)ds \right\} > 1.$$

If $(\varphi, \psi) \in K_1 \times K_1$, $\|(\varphi, \psi)\| = r$, and $T(\varphi, \psi) = \lambda(\varphi, \psi)$ for some $\lambda > 0$, then we get

$$\begin{aligned} \lambda(\varphi(t), \psi(t)) &= \left(\int_0^1 G(t, s)A(s)f(\varphi(s), \psi(s))ds, \right. \\ &\quad \left. \int_0^1 G(t, s)B(s)g(\varphi(s), \psi(s))ds \right). \end{aligned}$$

Picking $t_0 \in [\frac{1}{4}, \frac{3}{4}]$ such that

$$|(\varphi(t_0), \psi(t_0))| = \inf_{t \in [1/4, 3/4]} |(\varphi(t), \psi(t))| > 0,$$

then

$$\begin{aligned}
 & \lambda |(\varphi(t_0), \psi(t_0))| \\
 &= \inf_{t \in [1/4, 3/4]} \left| \left(\int_0^1 G(t, s) A(s) f(\varphi(s), \psi(s)) ds, \right. \right. \\
 & \quad \left. \left. \int_0^1 G(t, s) B(s) g(\varphi(s), \psi(s)) ds \right) \right| \\
 &\geq M \inf_{t \in [1/4, 3/4]} \left| \left(\int_0^1 G(t, s) A(s) |(\varphi(s), \psi(s))| ds, \right. \right. \\
 & \quad \left. \left. \int_0^1 G(t, s) B(s) |(\varphi(s), \psi(s))| ds \right) \right| \\
 &\geq M \inf_{t \in [1/4, 3/4]} \left| \left(\int_{1/4}^{3/4} G(t, s) A(s) \left| \left(\frac{\|\varphi\|}{4}, \frac{\|\psi\|}{4} \right) \right| ds, \right. \right. \\
 & \quad \left. \left. \int_{1/4}^{3/4} G(t, s) B(s) \left| \left(\frac{\|\varphi\|}{4}, \frac{\|\psi\|}{4} \right) \right| ds \right) \right| \\
 &= \frac{Mr}{4} \inf_{t \in [1/4, 3/4]} \left| \left(\int_{1/4}^{3/4} G(t, s) A(s) ds, \int_{1/4}^{3/4} G(t, s) B(s) ds \right) \right| \\
 &> r
 \end{aligned}$$

and, hence, $\lambda > 1$.

If $(H_4)(i)$ holds, say, $f(x, y) \leq N$ and $g(x, y) \leq N$, where N is a positive constant, then we choose

$$R > \max \left\{ 2r, N \int_0^1 G(t, s) A(s) ds, N \int_0^1 G(t, s) B(s) ds \right\}.$$

If $(\varphi, \psi) \in K_1 \times K_1$, $\|(\varphi, \psi)\| = R$, and

$$T(\varphi, \psi) = \lambda(\varphi, \psi) \quad \text{for some } \lambda > 0,$$

then there exists $t_0 \in [0, 1]$ such that

$$\|(\varphi, \psi)\| = |(\varphi(t_0), \psi(t_0))|$$

and, therefore,

$$\begin{aligned}
 \lambda |(\varphi(t_0), \psi(t_0))| &= \left| \left(\int_0^1 G(t_0, s) A(s) f(\varphi(s), \psi(s)) ds, \right. \right. \\
 & \quad \left. \left. \int_0^1 G(t_0, s) B(s) g(\varphi(s), \psi(s)) ds \right) \right| \\
 &< R;
 \end{aligned}$$

i.e., $\lambda < 1$.

If $(H_4)(ii)$ holds, then there is a $N > 0$ such that $g(x, y) \leq N$. From (2.3), there is a $R \in \Sigma_f$, $R > \max\{2r, N \int_0^1 G(s, s) B(s) ds\}$ such that

$$f(x, y) \leq \varepsilon |(x, y)| \quad \text{for } (x, y) \in \mathbb{R}^+ \times \mathbb{R}^+, |(x, y)| \geq R,$$

where the constant $\varepsilon > 0$ is such that

$$\varepsilon \int_0^1 G(s, s) A(s) ds < 1.$$

In particular, $|(x, y)| = R$ implies $f(x, y) \leq \varepsilon R$.

Let $(\varphi, \psi) \in K_1 \times K_1$, $\|(\varphi, \psi)\| = R$, and

$$T(\varphi, \psi) = \lambda(\varphi, \psi) \quad \text{for some } \lambda > 0.$$

Then there exists a $t_0 \in [0, 1]$ such that

$$\|(\varphi, \psi)\| = |(\varphi(t_0), \psi(t_0))|$$

and, therefore,

$$\begin{aligned} & \lambda |(\varphi(t_0), \psi(t_0))| \\ &= \left| \left(\int_0^1 G(t_0, s) A(s) f(\varphi(s), \psi(s)) ds, \right. \right. \\ & \quad \left. \left. \int_0^1 G(t_0, s) B(s) g(\varphi(s), \psi(s)) ds \right) \right| \\ &\leq \left| \left(\int_0^1 G(t_0, s) A(s) \max_{[0, R; 0, R]} f(x, y) ds, N \int_0^1 G(t_0, s) B(s) ds \right) \right| \\ &< R \end{aligned}$$

(we are able to do this since $R \in \Sigma_f$); i.e., $\lambda < 1$.

If $(H_4)(iii)$ holds, then there is $R \in \Sigma_f$, $R > 2r$ such that

$$\begin{aligned} f(x, y) &\leq \varepsilon |(x, y)|, \\ g(x, y) &\leq \varepsilon |(x, y)| \quad \text{for } (x, y) \in \mathbb{R}^+ \times \mathbb{R}^+, |(x, y)| \geq R, \end{aligned}$$

where the constant ε satisfies

$$\varepsilon \max \left\{ \int_0^1 G(s, s) A(s) ds, \int_0^1 G(s, s) B(s) ds \right\} < 1.$$

The monotonicity assumption on g implies that

$$R \in \Sigma_g$$

(cf. (3.5)).

Let $(\varphi, \psi) \in K_1 \times K_1$, $\|(\varphi, \psi)\| = R$, and

$$T(\varphi, \psi) = \lambda(\varphi, \psi) \quad \text{for some } \lambda > 0.$$

Then there is $t_0 \in [0, 1]$ such that

$$\|(\varphi, \psi)\| = |(\varphi(t_0), \psi(t_0))|$$

and, therefore,

$$\begin{aligned} \lambda |(\varphi(t_0), \psi(t_0))| &\leq \left| \left(\int_0^1 G(t_0, s) A(s) \max_{[0, R; 0, R]} f(x, y) ds, \right. \right. \\ &\quad \left. \left. \int_0^1 G(t_0, s) B(s) \max_{[0, R; 0, R]} g(x, y) ds \right) \right| \\ &\leq \varepsilon R \max \left\{ \int_0^1 G(s, s) A(s) ds, \int_0^1 G(s, s) B(s) ds \right\} \\ &< R; \end{aligned}$$

i.e., $\lambda < 1$.

If $(H_4)(iv)$ or $(H_4)(v)$ holds, a simple modification of the above argument yields that there exists $R > 2r$, such that

$$(\varphi, \psi) \in K_1 \times K_1, \quad \|(\varphi, \psi)\| = R, \quad T(\varphi, \psi) = \lambda(\varphi, \psi) \Rightarrow \lambda < 1.$$

The last step is to prove that

$$\inf_{\|(\varphi, \psi)\| = r} \|T(\varphi, \psi)\| > 0.$$

From (4.1), we get that for $\|(\varphi, \psi)\| = r$,

$$\begin{aligned} & \|T(\varphi, \psi)\| \\ & \geq \left\| \left(\int_0^1 G(t, s) A(s) M |(\varphi(s), \psi(s))| ds, \right. \right. \\ & \quad \left. \left. \int_0^1 G(t, s) B(s) M |(\varphi(s), \psi(s))| ds \right) \right\| \\ & \geq M \left\| \left(\int_{1/4}^{3/4} G(t, s) A(s) \left| \left(\frac{\|\varphi\|}{4}, \frac{\|\psi\|}{4} \right) \right| ds, \right. \right. \\ & \quad \left. \left. \int_{1/4}^{3/4} G(t, s) B(s) \left| \left(\frac{\|\varphi\|}{4}, \frac{\|\psi\|}{4} \right) \right| ds \right) \right\| \\ & \geq \frac{Mr}{4} \min \left\{ \int_{1/4}^{3/4} G(t, s) A(s) ds, \int_{1/4}^{3/4} G(t, s) B(s) ds \right\}, \end{aligned}$$

and the last expression is a positive number independent of (φ, ψ) .

Thus T has a fixed point $(\bar{\varphi}, \bar{\psi})$ in

$$D = \{(\varphi, \psi) \in K_1 \times K_1 \mid r \leq \|(\varphi, \psi)\| \leq R\}.$$

We claim that $\bar{\varphi}(t) > 0$, $\bar{\psi}(t) > 0$ for $t \in (0, 1)$. Indeed, from $\|(\bar{\varphi}, \bar{\psi})\| \geq r$ and Lemma 4, we know that $\bar{\varphi} > 0$ in $(0, 1)$ or $\bar{\psi} > 0$ in $(0, 1)$. We may assume that $\bar{\psi} > 0$ in $(0, 1)$. Again from $\|(\bar{\varphi}, \bar{\psi})\| > r$, there is $s_0 \in (0, 1)$ such that $\|(\bar{\varphi}(s_0), \bar{\psi}(s_0))\| = r$ and $\|(\bar{\varphi}(s), \bar{\psi}(s))\| \leq r$ for $0 < s \leq s_0$. This, together with (4.1) and Lemma 4, implies

$$\begin{aligned} f(\bar{\varphi}(s), \bar{\psi}(s)) & \geq M |(\bar{\varphi}(s), \bar{\psi}(s))| > 0, \\ g(\bar{\varphi}(s), \bar{\psi}(s)) & \geq M |(\bar{\varphi}(s), \bar{\psi}(s))| > 0, \quad 0 < s < s_0, \end{aligned}$$

and, hence,

$$\begin{aligned} \bar{\varphi}(t) & = \int_0^1 G(t, s) A(s) f(\bar{\varphi}(s), \bar{\psi}(s)) ds \\ & \geq \int_0^{s_0} G(t, s) A(s) f(\bar{\varphi}(s), \bar{\psi}(s)) ds > 0, \quad 0 < t < 1. \end{aligned}$$

Therefore (1.1)–(1.2)_a has a positive radial solution. This completes the proof of the first part of Theorem 1.

Next we consider (3.3)–(3.4)_b and (3.3)–(3.4)_c. It is easy to check that (3.3)–(3.4)_b and (3.3)–(3.4)_c are equivalent to the integral equations

$$(\varphi(t), \psi(t)) = \left(\int_0^1 K_1(t, s) A(s) f(\varphi(s), \psi(s)) ds, \right. \\ \left. \int_0^1 K_1(t, s) B(s) g(\varphi(s), \psi(s)) ds \right)$$

and

$$(\varphi(t), \psi(t)) = \left(\int_0^1 K_2(t, s) A(s) f(\varphi(s), \psi(s)) ds, \right. \\ \left. \int_0^1 K_2(t, s) B(s) g(\varphi(s), \psi(s)) ds \right),$$

respectively, where

$$K_1(t, s) = \begin{cases} t, & t \leq s, \\ s, & t > s, \end{cases} \quad K_2(t, s) = \begin{cases} 1-s, & t \leq s, \\ 1-t, & t > s. \end{cases}$$

For (3.3)–(3.4)_b let P_1 be the cone

$$P_1 = \left\{ z \in X \mid z(t) \geq 0, z(0) = z'(1) = 0, \min_{1/2 \leq t \leq 1} z(t) \geq \frac{\|z\|}{2} \right\};$$

for (3.3)–(3.4)_c let P_2 be the cone

$$P_2 = \left\{ z \in X \mid z(t) \geq 0, z'(0) = z(1) = 0, \min_{0 \leq t \leq 1/2} z(t) \geq \frac{\|z\|}{2} \right\}.$$

By the method above, the fixed-point theorem is used to prove that both (1.1)–(1.2)_b and (1.1)–(1.2)_c have positive solutions. This completes the proof of Theorem 1. ■

REFERENCES

1. C. Bandle, C. V. Coffman, and M. Marcus, Nonlinear elliptic problems in the annulus, *J. Differential Equations*, **69** (1987), 322–345.
2. K. Deimling, “Nonlinear Functional Analysis,” Springer, New York, 1985.

3. L. H. Erbe and H. Wang, On the existence of positive solutions of ordinary differential equations, *Proc. Amer. Math. Soc.* **120** (1994), 743–748.
4. J. A. Gatica and H. L. Smith, Fixed point theorems in cone with applications, *J. Math. Anal. Appl.* **61** (1977), 58–71.
5. P. L. Lions, On the existence of positive solutions of semilinear elliptic equations, *SIAM Rev.* **24** (1982), 441–467.
6. W. M. Ni and R. D. Nussbaum, Uniqueness and nonuniqueness for positive radial solutions of $\Delta u + f(x, r) = 0$, *Comm. Pure Appl. Math.* **38** (1985), 67–108.
7. H. Wang, On the existence of positive solutions for semilinear elliptic equations in the annulus, *J. Differential Equations* **109** (1994), 1–8.